

Acute Diarrhoea in Adults

Diarrhoea can be of sudden onset and lasting for less than four weeks (acute) or persistent (chronic). This leaflet deals with acute diarrhoea, which is common. In most cases, diarrhoea eases and goes within several days but sometimes takes longer. The main risk is lack of fluid in the body (dehydration). The main treatment is to have lots to drink, which aims to avoid dehydration. You should also eat as normally as possible. See a doctor if you suspect that you are dehydrating, or if you have any worrying symptoms such as those which are listed below.

What causes acute diarrhoea?

- **Infection** of the gut is the common cause. This is of sudden onset and is called acute infectious diarrhoea. Many bacteria, viruses and other germs can cause diarrhoea. Sometimes the germs come from infected food (food poisoning). Infected water is a cause in some countries. Sometimes it is just "one of those germs going about". Viruses are easily spread from one person to another by close contact, or when an infected person prepares food for others.
- **Other causes** are uncommon and include drinking lots of beer, side-effects from some medicines and anxiety.
- **Gut disorders** that cause persistent (chronic) diarrhoea may be mistaken for acute diarrhoea when they first begin - for example, diarrhoea caused by ulcerative colitis.

The rest of this leaflet deals only with infectious causes of acute diarrhoea. There are also other leaflets that give more details about some of the different germs (microbes) that cause infectious diarrhoea.

What are the symptoms of acute infectious diarrhoea?

- The main symptom is diarrhoea, often with being sick (vomiting) as well. Diarrhoea means loose or watery stools (faeces), usually at least three times in 24 hours. Blood or mucus can appear in the stools with some infections.
- Crampy pains in your tummy (abdomen) are common. Pains may ease for a while each time you pass some diarrhoea.
- A high temperature (fever), headache and aching limbs sometimes occur.

If vomiting occurs, it often lasts only a day or so but sometimes longer. Diarrhoea often continues after the vomiting stops and commonly lasts for several days or more. Slightly loose stools may persist for a week or so further before a normal pattern returns. Sometimes the symptoms last longer.

Symptoms of lack of fluid in the body (dehydration)

Diarrhoea and vomiting may cause dehydration. Consult a doctor quickly if you suspect you are becoming dehydrated. Mild dehydration is common and is usually easily and quickly reversed by drinking lots of fluids. Severe dehydration can be fatal unless quickly treated. This is because the organs of your body need a certain amount of fluid to function.

- Symptoms of dehydration in adults include:
 - Tiredness.
 - Dizziness or light-headedness.
 - Headache.
 - Muscular cramps.
 - Sunken eyes.
 - Passing little urine.
 - A dry mouth and tongue.
 - Weakness.

- Becoming irritable.
- Symptoms of *severe* dehydration in adults include:
 - Weakness.
 - Confusion.
 - Rapid heart rate.
 - Coma.
 - A greatly reduced amount of urine that you make.

This is a medical emergency and immediate medical attention is needed.

Dehydration in adults with acute diarrhoea is more likely to occur in:

- Elderly or frail people.
- Pregnant women.
- People with severe diarrhoea and vomiting. In particular, if you are not able to replace the fluid lost with enough drinks.

Do I need any tests?

Most people with acute infectious diarrhoea do not need to see a doctor or seek medical advice. Symptoms are often quite mild and commonly get better within a few days without any medical treatment.

However, in some circumstances, you may need to see a doctor (see below about when to seek medical advice). The doctor may ask you various questions - for example, about:

- Recent travel abroad.
- Whether you have been in contact with someone with similar symptoms.
- Whether you have recently taken antibiotics.
- Whether you have recently been admitted to hospital.

This is to look for possible causes of your diarrhoea. The doctor will also usually examine you, especially looking for signs of lack of fluid in the body (dehydration).

Tests are not usually needed. However, if you are particularly unwell, have bloody stools (faeces), have recently travelled abroad, are admitted to hospital, or your symptoms are not getting better, your doctor may ask you to collect a stool sample. This can then be examined in the laboratory to look for the cause of the infection.

When should I seek medical advice?

Seek medical advice in any of the following situations, or if any other symptoms occur that you are concerned about:

- If you suspect that you are becoming lacking in fluid in the body (dehydrated).
- If you are being sick (vomiting) a lot and unable to keep fluids down.
- If you have blood in your diarrhoea or vomit.
- If you have severe tummy (abdominal) pain.
- If you have severe symptoms, or if you feel that your condition is getting worse.
- If you have a persisting high temperature (fever).
- If your symptoms are not settling - for example, vomiting for more than 1-2 days, or diarrhoea that does not start to settle after 3-4 days.
- Infections caught abroad.
- If you are elderly or have an underlying health problem such as diabetes, epilepsy, inflammatory bowel disease, kidney disease.
- If you have a weakened immune system because of, for example, chemotherapy treatment, long-term steroid treatment, HIV infection.
- If you are pregnant.

What is the treatment for infectious diarrhoea in adults?

Symptoms often settle within a few days or so as your immune system usually clears the infection. Occasionally, admission to hospital is needed if symptoms are severe, or if complications develop (see below).

The following are commonly advised until symptoms ease.

Fluids - have lots to drink

The aim is to prevent lack of fluid in the body (dehydration), or to treat dehydration if it has developed. (**Note:** if you suspect that you are dehydrated, you should contact a doctor.)

- As a rough guide, drink at least 200 mls after each bout of diarrhoea (after each watery stool (faeces)).
- This extra fluid is in addition to what you would normally drink. For example, an adult will normally drink about two litres a day but more in hot countries. The above advice of 200 mls after each bout of diarrhoea is in addition to this usual amount that you would drink.
- If you have been sick (vomited), wait 5-10 minutes and then start drinking again but more slowly. For example, a sip every 2-3 minutes but making sure that your total intake is as described above.
- You will need to drink even more if you are dehydrated. A doctor will advise on how much to drink if you are dehydrated.

For most adults, fluids drunk to keep hydrated should mainly be water. It is best not to have drinks that contain a lot of sugar, such as cola or pop, as they can sometimes make diarrhoea worse.

Rehydration drinks are recommended for people who are frail, or over the age of 60, or who have underlying health problems. They are made from sachets that you can buy from pharmacies. (The sachets are also available on prescription.) You add the contents of the sachet to water. Rehydration drinks provide a good balance of water, salts and sugar. They do not stop or reduce diarrhoea. However, the small amount of sugar and salt helps the water to be absorbed better from the gut into the body. Home-made salt/sugar mixtures are used in developing countries if rehydration drinks are not available but they have to be made carefully, as too much salt can be dangerous. Rehydration drinks are cheap and readily available in the UK and are the best treatment.

Anti-secretory medicines are designed to be used with rehydration treatment. They reduce the amount of water that is released into the gut during an episode of diarrhoea. They can be used for children who are older than 3 months of age, and for adults.

Eat as normally as possible

It used to be advised to not eat for a while if you had infectious diarrhoea. However, now it is advised to eat small, light meals if you can. Be guided by your appetite. You may not feel like food and most adults can do without food for a few days. Eat as soon as you are able - but don't stop drinking. If you do feel like eating, avoid fatty, spicy or heavy food at first. Plain foods such as wholemeal bread and rice are good foods to try eating first.

Medication

Antidiarrhoeal medicines are not usually necessary. However, you may wish to reduce the number of trips that you need to make to the toilet. You can buy antidiarrhoeal medicines from pharmacies. The safest and most effective is **loperamide**. The adult dose of this is two capsules at first. This is followed by one capsule after each time you pass some diarrhoea, up to a maximum of eight capsules in 24 hours. It works by slowing down your gut's activity. You should not take loperamide for longer than five days.

Note: do not give antidiarrhoeal medicines to children aged under 12 years. Also, do not use antidiarrhoeal medicines if you pass blood or mucus with the diarrhoea or if you have a high temperature (fever). People with certain conditions should not take loperamide. Therefore, read the leaflet that comes with the medicine to be safe. For example, pregnant women should not take loperamide.

Paracetamol or **ibuprofen** are useful to ease a high temperature or headache.

As explained above, if symptoms are severe, or continue (persist) for more than several days, your doctor may ask for a sample of the diarrhoea. This is sent to the laboratory to look for infecting germs (bacteria, parasites, etc). Sometimes an antibiotic or other treatments are needed if certain bacteria or other infections are found to be the cause. Antibiotics are of no use for infectious diarrhoea caused by viruses and may even make things worse.

Are there any complications that may occur?

Complications are uncommon in the UK. They are more likely in the very young, in pregnant women, or in the elderly. They are also more likely if you have an ongoing (chronic) disease such as diabetes, or if your immune system may not be working fully. For example, if you are taking long-term steroid medication or you are having chemotherapy treatment for cancer.

Possible complications include the following:

- **Lack of fluid (dehydration) and salt (electrolyte) imbalance in your body.** This is the most common complication. It occurs if the water and salts that are lost in your stools (faeces), or when you have been sick (vomited), are not replaced by your drinking enough fluids. If you can manage to drink plenty of fluids then dehydration is unlikely to occur, or is only likely to be mild and will soon recover as you drink. Severe dehydration can lead to a drop in your blood pressure. This can cause reduced blood flow to your vital organs. If dehydration is not treated, kidney failure may also develop. Some people who become severely dehydrated need a drip of fluid directly into a vein. This requires admission to hospital.
- **Reactive complications.** Rarely, other parts of the body may react to an infection that occurs in the gut. This can cause symptoms such as **joint inflammation (arthritis)**, skin inflammation and eye inflammation (either conjunctivitis or **uveitis**). Reactive complications are uncommon if you have a virus causing infectious diarrhoea.
- **Spread of infection** to other parts of your body such as your bones, joints, or the meninges that surround your brain and spinal cord. This is rare. If it does occur, it is more likely if **diarrhoea is caused by *Salmonella* spp. infection.**
- **Persistent syndromes of diarrhoea** may rarely develop.
- **Irritable bowel syndrome** is sometimes triggered by a bout of infectious diarrhoea.
- **Lactose intolerance can sometimes occur** for a period of time after infectious diarrhoea. This is known as secondary or acquired lactose intolerance. Your gut lining can be damaged by the episode of diarrhoea. This leads to lack of an enzyme (chemical) called lactase that is needed to help your body digest a sugar called lactose that is in milk. Lactose intolerance leads to bloating, tummy (abdominal) pain, wind and watery stools after drinking milk. The condition gets better when the infection is over and the gut lining heals. It is more common in children.
- **Haemolytic uraemic syndrome** is another potential complication. It is rare and is usually associated with infectious **diarrhoea caused by a certain type of *Escherichia coli* infection.** It is a serious condition where there is anaemia, a low platelet count in the blood and kidney failure. It is more common in children. If recognised and treated, most people recover well.
- **Reduced effectiveness of some medicines.** During an episode of infectious diarrhoea, certain medicines that you may be taking for other conditions or reasons may not be as effective. This is because the diarrhoea and/or vomiting means that reduced amounts of the medicines are taken up (absorbed) into your body. Examples of such medicines are **medicines for epilepsy, diabetes and contraception.** Speak to your doctor or practice nurse if you are unsure of what to do if you are taking other medicines and have acute diarrhoea.

Preventing spread of infection to others

Some infections causing diarrhoea are very easily passed on from person to person. If you have acute diarrhoea, the following are also recommended to prevent the spread of infection to others:

- Wash your hands thoroughly after going to the toilet. Ideally, use liquid soap in warm running water but any soap is better than none. Dry properly after washing.
- Don't share towels and flannels.
- Don't prepare or serve food for others.
- Regularly clean the toilets that you use. Wipe the flush handle, toilet seat, bathroom taps, surfaces and door handles with hot water and detergent at least once a day. Keep a cloth just for cleaning the toilet (or use a disposable one each time).

- Stay off work, college, etc, until at least 48 hours after the last episode of diarrhoea or being sick (vomiting).
- Food handlers: if you work with food and develop diarrhoea or vomiting, you must immediately leave the food-handling area. For most, no other measures are needed, other than staying away from work until at least 48 hours after the last episode of diarrhoea or vomiting. Some special situations may arise and sometimes longer time off is needed. Specialist advice may be needed for some uncommon causes of infectious diarrhoea. If in doubt, seek advice from your employer or GP.
- If the cause of acute diarrhoea is known to be (or suspected to be) a germ called *Cryptosporidium* spp., you should not swim in swimming pools for two weeks after the last episode of diarrhoea.

Can infectious diarrhoea be prevented?

The advice given in the previous section is mainly aimed at preventing the spread of infection to other people. However, even when we are not in contact with someone with infectious diarrhoea, proper storage, preparation and cooking of food and good hygiene help to prevent it.

In particular, always wash your hands:

- After you go to the toilet.
- Before you touch food.
- Between handling raw meat and food ready to be eaten. (There may be some germs (bacteria) on raw meat.)
- After gardening.
- After playing with pets (healthy animals can carry certain harmful bacteria).

The simple measure of washing hands regularly and properly is known to make a big difference to the chance of developing infectious diarrhoea.

You should also take extra measures when in countries of poor sanitation. For example, avoid water and other drinks that may not be safe and avoid food washed in unsafe water.

Further reading & references

- Ejemot RI, Ehiri JE, Meremikwu MM, et al; Hand washing for preventing diarrhoea. *Cochrane Database Syst Rev*. 2008 Jan 23;(1):CD004265.
- *Acute diarrhoea in adults: racecadotril*; NICE Evidence Summary *New Medicine*, March 2013
- *Diarrhoea and vomiting in children under 5*; NICE Clinical Guideline (April 2009)
- *Preventing person-to-person spread following gastrointestinal infections: guidelines for public health physicians and environmental health officers*; GOV.UK (2005)

Disclaimer: This article is for information only and should not be used for the diagnosis or treatment of medical conditions. EMIS has used all reasonable care in compiling the information but make no warranty as to its accuracy. Consult a doctor or other health care professional for diagnosis and treatment of medical conditions. For details see our [conditions](#).

Original Author: Dr Tim Kenny	Current Version: Dr Laurence Knott	Peer Reviewer: Prof Cathy Jackson
Document ID: 4521 (v43)	Last Checked: 01/12/2014	Next Review: 30/11/2017

View this article online at www.patient.co.uk/health/acute-diarrhoea-in-adults-leaflet.

Discuss Acute Diarrhoea in Adults and find more trusted resources at www.patient.co.uk.

© EMIS. EMIS is a trading name of Egton Medical Information Systems Limited.